

Exponential and Index Functions

Index laws

$$\begin{aligned}
 a^m \times a^n &= a^{m+n} \\
 a^m \div a^n &= a^{m-n} \\
 (a^m)^n &= a^{m \cdot n} \\
 (ab)^m &= a^m b^m \\
 \left(\frac{a}{b}\right)^m &= \frac{a^m}{b^m}
 \end{aligned}$$

Fractional indices

$${}^n\sqrt{x} = x^{1/n}$$

Logarithms

$$\log_b(x) = n \quad \text{where } b^n = x$$

Using logs to solve index eq's

Used for equations without common base exponent

Or change base:

$$\log_b c = \frac{\log_a c}{\log_a b}$$

If $a < 1$, $\log_b a < 0$ (flip inequality operator)

Exponential functions

e^x - natural exponential function

$$\lim_{h \rightarrow 0} \frac{e^h - 1}{h} = 1$$

Logarithm laws

$$\begin{aligned}
 \log_a(mn) &= \log_a m + \log_a n \\
 \log_a\left(\frac{m}{n}\right) &= \log_a m - \log_a n \\
 \log_a(m^p) &= p \log_a m \\
 \log_a(m^{-1}) &= -\log_a m \\
 \log_a 1 &= 0 \quad \text{and} \quad \log_a a = 1
 \end{aligned}$$

Inverse functions

For $f: \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = a^x$, inverse is:

$$f^{-1}: \mathbb{R}^+ \rightarrow \mathbb{R}, f^{-1} = \log_a x$$

Euler's number

$$e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$$

Exponential and logarithmic modelling

$$A = A_0 e^{kt}$$

where

A_0 is initial value

t is time taken

(1) k is a constant


For continuous growth, $k > 0$

For continuous decay, $k < 0$

Graphing exponential functions

$$f(x) = Aa^{k(x-b)} + c, \quad |a > 1$$

- **y-intercept** at $(0, A \cdot a^{-kb} + c)$ as $x \rightarrow \infty$
- **horizontal asymptote** at $y = c$
- **domain** is \mathbb{R}
- **range** is (c, ∞)
- dilation of factor A from x -axis
- dilation of factor $\frac{1}{k}$ from y -axis




Graphing logarithmic functions

$\log_e x$ is the inverse of e^x (reflection across $y = x$)

$$f(x) = A \log_a k(x - b) + c$$

where

- **domain** is (b, ∞)
- **range** is \mathbb{R}
- **vertical asymptote** at $x = b$
- **y-intercept** exists if $b < 0$
- dilation of factor A from x -axis
- dilation of factor $\frac{1}{k}$ from y -axis



Finding equations

Solve simultaneous equations on CAS: $\begin{cases} f(3)=9 \\ g(3)=0 \end{cases} \Big|_{a,b}$