


Circular functions

Exact values



$$1 \text{ rad} = \frac{180 \text{ deg}}{\pi}$$

sin and cos graphs

$$f(x) = a \sin(bx - c) + d$$

$$f(x) = a \cos(bx - c) + d$$

where

- a is the y -dilation (amplitude)
- b is the x -dilation (period)
- c is the x -shift (phase)
- d is the y -shift (equilibrium position)

Domain is \mathbb{R}

Range is $[-b + c, b + c]$;

Graph of $\cos(x)$ starts at $(0, 1)$. Graph of $\sin(x)$ starts at $(0, 0)$.

Mean / equilibrium: line that the graph oscillates around ($y = d$)

Amplitude

Graph oscillates between $+a$ and $-a$ in y -axis

$a = 0$ produces straight line

$a < 0$ inverts the phase (sin becomes cos, vice versa)

Period

Period T is $\frac{2\pi}{b}$

$b = 0$ produces straight line

$b < 0$ inverts the phase

Phase

c moves the graph left-right in the x axis.

If $c = T = \frac{2\pi}{b}$, the graph has no actual phase shift.

Symmetry

$$\sin\left(\theta + \frac{\pi}{2}\right) = \cos \theta$$

$$\sin(\theta + \pi) = -\sin \theta$$

$$\cos\left(\theta + \frac{\pi}{2}\right) = -\sin \theta$$

$$\cos(\theta + \pi) = -\cos\left(\theta + \frac{3\pi}{2}\right) = \cos(-\theta)$$

Pythagorean identity

$$\cos^2 \theta + \sin^2 \theta = 1$$

Complementary relationships

$$\sin\left(\frac{\pi}{2} - \theta\right) = \cos \theta$$

$$\cos\left(\frac{\pi}{2} - \theta\right) = \sin \theta$$

$$\sin \theta = -\cos\left(\theta + \frac{\pi}{2}\right)$$

$$\cos \theta = \sin\left(\theta + \frac{\pi}{2}\right)$$

tan graph

$$y = a \tan(nx)$$

where

- a is x -dilation (period)
- n is y -dilation (\equiv amplitude)
- period T is $\frac{\pi}{n}$
- range is \mathbb{R}
- roots at $x = \frac{k\pi}{n}$
- asymptotes at $x = \frac{(2k+1)\pi}{2n}$, $k \in \mathbb{Z}$

Asymptotes should always have equations and arrow pointing up

Solving trig equations

1. Solve domain for $n\theta$
2. Find solutions for $n\theta$
3. Divide solutions by n

$$\sin 2\theta = \frac{\sqrt{3}}{2}, \quad \theta \in [0, 2\pi] \quad (\because 2\theta \in [0, 4\pi])$$

$$2\theta = \sin^{-1} \frac{\sqrt{3}}{2}$$

$$2\theta = \frac{\pi}{3}, \frac{2\pi}{3}, \frac{7\pi}{3}, \frac{8\pi}{3}$$

$$\therefore \theta = \frac{\pi}{6}, \frac{\pi}{3}, \frac{7\pi}{6}, \frac{4\pi}{3}$$