

Polynomials

Quadratics

General form	$x^2 + bx + c = (x + m)(x + n)$ where $mn = c, m + n = b$
--------------	--

Difference of squares	$a^2 - b^2 = (a - b)(a + b)$
-----------------------	------------------------------

Perfect squares	$a^2 \pm 2ab + b^2 = (a \pm b)^2$
-----------------	-----------------------------------

Completing the square	$x^2 + bx + c = (x + \frac{b}{2})^2 + c - \frac{b^2}{4}$ $ax^2 + bx + c = a(x - \frac{b}{2a})^2 + c - \frac{b^2}{4a}$
-----------------------	--

Quadratic formula	$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ where $\Delta = b^2 - 4ac$
-------------------	---

Cubics

Difference of cubes: $a^3 - b^3 = (a - b)(a^2 + ab + b^2)$

Sum of cubes: $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$

Perfect cubes: $a^3 \pm 3a^2b + 3ab^2 \pm b^3 = (a \pm b)^3$

$$y = a(bx - h)^3 + c$$

- $m = 0$ at *stationary point of inflection* (i.e. $(\frac{h}{b}, k)$)
- in form $y = (x - a)^2(x - b)$, local max at $x = a$, local min at $x = b$
- in form $y = a(x - b)(x - c)(x - d)$: x -intercepts at b, c, d
- in form $y = a(x - b)^2(x - c)$, touches x -axis at b , intercept at c

Linear and quadratic graphs

Forms of linear equations

$y = mx + c$ where m is gradient and c is y -intercept
 $\frac{x}{a} + \frac{y}{b} = 1$ where m is gradient and (x_1, y_1) lies on the graph

$y - y_1 = m(x - x_1)$ where $(a, 0)$ and $(0, b)$ are x - and y -intercepts

Line properties

Parallel lines: $m_1 = m_2$

Perpendicular lines: $m_1 \times m_2 = -1$

Distance: $|\vec{AB}| = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

Quartic graphs

Forms of quadratic equations

$$y = ax^4$$

$$y = a(x - b)(x - c)(x - d)(x - e)$$

$$y = ax^4 + cd^2 (c \geq 0)$$

$$y = ax^2(x - b)(x - c)$$

$$y = a(x - b)^2(x - c)^2$$

$$y = a(x - b)(x - c)^3$$

Simultaneous equations (linear)

- Unique solution** - lines intersect at point
- Infinitely many solutions** - lines are equal
- No solution** - lines are parallel

Solving $\begin{cases} px + qy = a \\ rx + sy = b \end{cases}$ for $\{0, 1, \infty\}$ solutions

where all coefficients are known except for one, and a, b are known

- Write as matrices: $\begin{bmatrix} p & q \\ r & s \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} a \\ b \end{bmatrix}$
- Find determinant of first matrix: $\Delta = ps - qr$
- Let $\Delta = 0$ for number of solutions $\neq 1$ or let $\Delta \neq 0$ for one unique solution.
- Solve determinant equation to find variable
 - for infinite/no solutions: —
- Substitute variable into both original equations
- Rearrange equations so that LHS of each is the same
- RHS(1) = RHS(2) $\implies (1) = (2) \forall x$ (∞ solns)
 RHS(1) \neq RHS(2) $\implies (1) \neq (2) \forall x$ (0 solns)

On CAS: Matrix \rightarrow det

Solving $\begin{cases} a_1x + b_1y + c_1z = d_1 \\ a_2x + b_2y + c_2z = d_2 \\ a_3x + b_3y + c_3z = d_3 \end{cases}$

- Use elimination
- Generate two new equations with only two variables
- Rearrange & solve
- Substitute one variable into another equation to find another variable
- etc.